

State Water Resources Control Board
Hearing Name IID Transfer - Phase 2

Exhibit: *A*

For Ident: _____ In Evidence: _____

**Synthesis Document
of Current Information on the
Sediment Physical Characteristics and Contaminants
at the Salton Sea
Riverside and Imperial Counties, California**

**March 3, 1999
LFR 6824.00-01**

**Prepared for
Salton Sea Authority
78-035 Calle Estado
La Quinta, California 92253-2930**

 **LFR**
LEVINE • FRICKE

 Printed on recycled materials

CONTENTS

1.0 INTRODUCTION.....	1
1.1 Background.....	1
2.0 SEDIMENTS AND SEDIMENTATION.....	2
3.0 CHEMICAL DATA	4
4.0 CONCLUSIONS.....	5
5.0 REFERENCES.....	7

TABLES

- 1 Concentrations of Inorganic Chemicals in Sediment from the Salton Sea and Surrounding Tributaries Determined To Be of Concern
- 2 Concentrations of Organic Chemicals in Sediment from the Salton Sea and Surrounding Tributaries Determined To Be of Concern

FIGURES

- 1 Salton Sea Bottom Sediment Sampling Sites

1.0 INTRODUCTION

LFR Levine-Fricke (LFR) has completed a review of records contained within the archives of the University of Redlands Salton Sea Database Program (SSDP). A list of the reference documents requested and reviewed from the SSDP is presented in Section 5.0 of this report. For some of the references, only abstracts were available. References not received or that did not pertain to the Salton Sea ("the Sea") bottom sediment characteristics are not listed. All of the information presented below was obtained from the references listed.

1.1 Background

Extensive research has been performed in and around the Sea to characterize water quality and to evaluate biological impacts from contaminants. Even more geotechnical research has been conducted on the tectonics and geologic setting of the Salton Trough. However, information on the bottom sediment characteristics and contaminants of the Salton Sea is limited.

Previous studies on the Salton Sea bottom sediments have identified a variety of inorganic and organic chemicals including organochlorine pesticide residues of banned DDT [1,1,1-trichloro-2,2-bis (p-chlorophenyl)-ethane] and its derivatives, DDD [1,1-dichloro-2,2-bis (p-chlorophenyl)-ethane], and DDE [1,1-dichloro-2,2-bis (p-chlorophenyl) ethylene]. Many of these same chemicals, plus some additional ones, have been identified in the riverbeds feeding into the Sea, including DDT, DDD, DDE, dichloromethane, polychlorinated biphenyls (PCBs), polynuclear aromatic hydrocarbons (PAHs), pesticides, selenium, and boron. Little is known about the current concentrations of these contaminants in sediments throughout the entire Salton Sea. Limited chemical data were collected by Bechtel (1997) at the Salton Sea Test Base, which comprises 13,462 acres of water located along Highway 86 at the southwest corner of the Sea and approximately 6 miles south of Salton City, and from Setmire and Stroud's (1990) irrigation study of the deltas and tributaries of the New and Alamo Rivers. Other documents which have provided the best available information to date on bottom sediment contaminants include Bechtel (1997), Eccles (1979), Hogg (1963), and Setmire [et al] (1993).

The geologic setting of the Salton Sea occurs in an area known as the Salton Trough. Some of the major contributors to the understanding of the tectonic systems present within the Salton Trough include Moran (1977), Babcock (1974), Thornton and Seyfried (1975), and Johnson et al (1994). Babcock (1974) noted that prior to deposition of the Borrego Formation, beds of the Shavers Well Formation were tilted and eroded, creating an unconformity upon which the Borrego lacustrine sediments were deposited and resulting in the current formation of the Salton Trough. It underwent repeated periods of desiccation interspersed with influxes of clayey/silty sediments, largely derived from the Colorado River. Thornton and Seyfried (1975)

noted that the sediments of the Salton Trough and the Gulf Coast region contain detritus and organic matter.

The report prepared by Bechtel National, Inc. for the U.S. Navy's removal site evaluation provided information regarding historic use of the Salton Sea Test Base property. The aeroballistic marine target area was reportedly used for testing inert atomic weapons. Approximately 3,750 test units were dropped into the Sea. These inert (non-explosive, non-radioactive) test units were usually stainless steel casings filled with arming, fusing, and firing components (containing lead/acid batteries [until the 1950s] or nickel/cadmium batteries, minor amounts of aluminum, copper, brass, and rubber), as well as concrete, lead and/or stainless steel ballast. Most units weighed between 5,000 and 40,000 pounds. Lead/acid and nickel/cadmium batteries were used for firing/fusing tests. The test units reportedly broke apart upon impact, scattering debris across the Sea floor. Approximately 10,000 pounds of material was recovered from one area investigated, but the majority of the debris still lies buried in the sediment of the Sea. A MK-6 "fly-around" radioactive test unit that contained 120 pounds of uranium was also reportedly lost in the Sea.

2.0 SEDIMENTS AND SEDIMENTATION

The Bechtel (1997) report also provided extensive sediment core data for the area of the Sea within the Navy's offshore property. Approximately 107 sediment samples were collected at three main areas of investigation. Core depths varied from 1 to 3 feet. These offshore areas included a shoreline disposal area (25 sediment samples), an offshore aeroballistic marine target area (68 sediment samples), and an Imhoff Tank area (14 sediment samples). Sediments encountered during these investigations consisted of predominantly sands to depths greater than 3 feet within 500 feet offshore, predominantly clay to depths greater than 3 feet between 500 and 12,000 feet offshore, and predominantly clay to a depth of 2.75 feet underlain by sand beyond 12,000 feet offshore.

In 1996, James Walker conducted sediment sampling at 59 sites using a Wildco coring device that measured 5 cm in diameter and 51 cm long (Walker, 1996). One sampling transect was conducted along a north-south baseline, and five sampling transects were conducted along an east-west baseline. Although the cores were shallow and no chemical analyses were run, these cores provide a representation of the grain sizes found in the bottom sediment. A summary of the grain size distribution for bottom sediments from this study is presented on Figure 1.

Ferrari and Weghorst (1997) completed a detailed survey to develop underwater topography, compute area-capacity relationships, and develop detailed bathymetry for design analysis. The bathymetric survey was run using sonic depth recording equipment interface with DGPS (accuracy of 1 - 2 meters), creating an above average quality map of the Sea.

In 1968, Van de Kamp (1973) investigated all the major facies within outcrops of the entire basin including lacustrine deposits, meandering channel deposits, alluvial fans and braided-stream deposits, and Aeolian sand deposits. Although none of the 18 core borings were collected from the Salton Sea bottom sediment, these cores provide information about the distribution and sources of sand and sediments within the Salton Sea watershed. The two major sources of sediments identified include the Colorado River and the basin margins. The Colorado River carried eroded debris from the Colorado Plateau to the southern part of the basin, depositing sand and mud in deltaic and lacustrine facies. The sediment deposits from sources at the basin margins were deposited in alluvial fans, braided streams, barrier beaches, and lacustrine beds.

Stephen (1972) investigated the New River delta and found it to have an extent greater than 15 km², draining 6,500 km² over its 150 km length. The suspended sediment load carried within the New River was estimated to be approximately 5.0×10^8 kg per year. This document provides descriptions of sediment mineralogy and grain size, and elaborates on the correlation of sediment size distribution with distributary patterns. Three years later, Stephen et al (1975) revisited the New River and reported on subaerial deposits comprising distributary channel, levee, and interdistributary subaerial flat and crevasse deposits. The investigators identified subaqueous deposits as largely prodelta clay and delta-front fine silt.

General conditions that affect sediments and distribution were best presented in Arnal (1961). The Sea's lowest elevation is -276.7 feet below sea level, with an annual temperature range from 10 to 34.5 Celsius. The Sea's currents move in a counterclockwise, gyral motion around the lake due to the influence of prevailing winds. Sand, silts, and clays are deposited in that order from the shore toward the center of the lake, where more fine sediments accumulate. The water content, amount of calcium carbonate, and natural characteristics indicate that most of the sediments (75 percent) were derived from the suspended load of the Colorado River, whereas the mineralogy suggests that some of the sediments have a local origin. The water content of the sediments varies in inverse ratio to the grain size, high (> 50 percent) in clay depositions (the deepest lake sediment; grain size less than 4 microns) and low (~ 20 percent) where sand is deposited. The water content decreases with depth. The pH of the sediments is regulated by a variety of physical and chemical properties and reactions including carbonates, organic matter, carbon dioxide, and organic acids from the decomposition of plant and animal matter. The distribution of the organic content of the sediments is influenced by phytoplankton, the texture of the sediments, and currents. The distribution shows a low organic content (< 1 percent) along the shore, with higher values (4 - 6 percent) found in the central part, and a maximum content (> 6 percent) found 3 miles offshore, near Fish Spring. In all sediments, quartz and plagioclases are the dominant primary minerals.

Inflow rates to the Sea were calculated using limited suspended sampling data and historical suspended sediment sampling data observed from other major reservoirs in the southwestern United States (United States Department of the Interior, 1970). The Salton Sea watershed is approximately 8,360 square miles. The long-term average sediment inflow volume calculated was 4,000 acre-feet of sediment per year. Over a

50-year period, this would amount to less than 4 percent of the gross water storage volume of the Salton Sea. A quarterly sediment sampling program carried out by the Imperial Irrigation District since 1952 reported a flow rate ranging from 1,508 to 385 cubic feet per second (cfs) for the Alamo River, and 964 to 315 cfs for the New River. The total average annual sediment contribution from the Alamo River and New River is estimated to be 340 acre-feet and 370 acre-feet, respectively. (Both estimates include an average flow rate and a 10 percent bedload pickup.) Sedimentation rates from 1928 through 1958 estimated at least 5 feet of sediments being deposited within the deltas of the Alamo and New Rivers. The sedimentation rate at the deltas was estimated to be 2 inches per year, as opposed to a rate of 0.02 inches per year for the central part of the lake (Arnal, 1961).

3.0 CHEMICAL DATA

Setmire & Stroud's (1990) document, which focused on agricultural runoff and drainage, measured sediment concentrations of "trace elements" (i.e., inorganic compounds) and organochlorine compounds in and around the Salton Sea. Seventeen sediment samples were collected from the upper 5 to 10 cm of sediments in 1986 and analyzed for "trace elements," organochlorine pesticides, and (to a limited extent) organophosphorous pesticides. The data are summarized in Table 1. The laboratory procedures document that samples were air dried, pulverized with a mortar and pestle, then split and sieved separately to eliminate sand particles that did not pass a 100-mesh and 230-mesh sieve (Stephen, 1992). Data were reported in dry weight. Chemicals detected above the maximum "baseline value" for soils of the western United States (Shacklette and Boerngen, 1984) were identified as being of concern. Setmire & Stroud concluded that the following chemicals were of concern: chromium, nickel, selenium, thorium, uranium, and zinc. Silver and cadmium were not detected in the bottom sediment samples. Median concentrations of Setmire & Stroud's "trace elements" were reported: 5.6 mg/kg arsenic, 0.7 mg/kg selenium, <2 mg/kg silver, 550 mg/kg barium, <2 mg/kg cadmium, 58 mg/kg chromium, 28 mg/kg copper, 21 mg/kg lead, <2 mg/kg molybdenum, 25 mg/kg nickel, 77 mg/kg vanadium, 78 mg/kg zinc, 10.6 mg/kg thorium, and 4.9 mg/kg uranium.

In this same 1990 Setmire and Stroud study, pesticides and PCBs were detected in bottom sediments. Other chemicals found include Chlordane in the New River at the international boundary, Toxaphene at Trifolium Drain 4, and Methoxychlor at Vail Drain 4. Table 2 summarizes these findings.

Five years later, Setmire et al (1993) published another study that focused more on surface and subsurface water quality and on biotic tissue concentrations of "trace elements," heavy metals, and organochlorine pesticides (Setmire et al, 1993). Of particular interest to the sedimentology of the Sea, this study investigated a naturally occurring "selenium removal process" at the mouth of the Alamo River. In August 1988 and February 1989, 16 samples of the bottom sediment were collected at the Alamo River delta and analyzed for selenium content. Selenium concentrations were between 0.2 and 0.3 mg/kg in the river sediment samples, and varied from 0.2 to

2.5 mg/kg at sites throughout the Alamo River delta. Relatively high levels of selenium (1.3 to 2.5 mg/kg) were found in the embayments, without any discernable pattern of distribution. This area of investigation is depicted in Figure 1. Analytical results are summarized in Table 2.

The Bechtel (1997) report stated that organochlorine pesticides, PAHs, and volatile organic compounds (acetone, carbon disulfide, ethylbenzene, toluene, and xylenes) were detected in sediment samples collected from the shoreline disposal area. Elevated concentrations of copper (68.7 mg/kg), barium, and thallium were also detected.

Elevated concentrations of cadmium (maximum 1.6 mg/kg), arsenic (maximum 27.4 mg/kg), antimony (maximum 9.9 mg/kg), molybdenum (maximum 14.5 mg/kg), selenium (maximum 8.4 mg/kg), and vanadium (maximum 52.5 mg/kg) were detected in the offshore aeroballistic marine target area sediment. A localized area of elevated uranium (maximum 14.2 mg/kg) was also identified. The report concluded that: 1) these contaminants were naturally occurring, with the exception of cadmium; and 2) based on the limited source and nature of the cadmium release (nickel/cadmium battery), no further action was warranted.

Organochlorine pesticides (DDE, Dieldrin, gamma-Chlordane, and/or Heptachlor) were detected in 3 of the 14 sediment samples collected from the Imhoff Tank area. Phenol was also detected in one of the samples, and thallium (maximum 0.26 mg/kg) was detected in two samples. Bechtel concluded that except for the organochlorine pesticides (attributed to irrigation drainage), the presence of these contaminants in sediment did not present a significant risk. It is unclear if the data presented in Bechtel (1997) were reported in dry or wet weights. Table 2 summarizes these findings.

Hogg (1973) performed some of the earliest pesticide work on the bottom sediments. He collected 6 substratum samples using SCUBA gear and 16mm (inner diameter) by 23cm long coring tubes. Mean values (reported in micrograms per kilogram; $\mu\text{g}/\text{kg}$) for pesticide residues of Dieldrin, DDT, DDD, DDE, and combined samples for the upper and lower layers of the core samples are summarized in Table 2. Based on his small data set for sediment contaminants, Hogg calculated the presence of 10,400 pounds of total DDT and its metabolites in the upper 12 cm for the entire Sea.

Eccles (1979) provided values for DDE concentrations in bottom sediment samples collected in tributaries to the Salton Sea. Eccles collected samples in 1977 and found concentrations of DDE at Avenue 64 Evacuation Channel (67 $\mu\text{g}/\text{kg}$) and at Trifolium Drain 1 (110 $\mu\text{g}/\text{kg}$).

4.0 CONCLUSIONS

Limited data representing current conditions of the Salton Sea bottom sediment types and contaminants are available. However, sufficient data have been collected over the past 26 years to show that the Salton Sea and surrounding tributary bottom sediments

include a variety of metals, metalloids, radioactive elements, pesticides, and organic compounds. To date, little is known about the sediment types in the Sea below depths of approximately 91 cm. Data regarding current concentrations of sediment contaminants in the Sea are also quite limited.

5.0 REFERENCES

- Arnal, Robert E. 1954. Preliminary Report on the Sediments and Foraminifera from the Salton Sea, Southern California [Abstracts]. Geological Society of America Bulletin. Vol. 65, Issue 12, p 2, pp. 1227-1228.
- Arnal, Robert E. 1961. Limnology, Sedimentation, and Microorganisms of the Salton Sea, California. Geological Society of America Bulletin. Vol. 72, Issue 3, pp. 427-78.
- Babcock, E. A. 1974. Geology of northeast margin of the Salton Trough, Salton Sea, California Geological Society of America Bulletin. Volume 85, Issue 3, pp. 321-332.
- Bechtel National, Inc. 1997. Final Addendum to the Removal Site Evaluation Report, Salton Sea Test Base Report, Salton Sea Test Base, Imperial County, CA. I v.
- Eccles, L.A. 1979. Pesticide residues in agricultural drains, southeastern desert area, California. U.S. Geological Survey Water Resources Investigation Report 79-16, 60 pp.
- Ferrari, R. L., and P. Weghorst. 1997. Salton Sea: 1995 Hydrographic GPS Survey (Revised May 1997). NTIS Accession # PB97 - 164677, p. 30.
- Hogg, N. D. 1973. Chlorinated Hydrocarbon Pesticide Residues, Salton Sea, California. M.S. Thesis. California Polytechnic University, Pomona. 47 pp.
- Johnson, H.; Agnew, D.; and Wyatt, F. 1994. Present-day crustal deformation in southern California. Journal of Geophysical Research. Vol. 99, Issue B12, pp. 23951- 23974.
- Morton, P. 1977. Geology and mineral resources of Imperial County, California. Sacramento, CA. California Division of Mines & Geology. p. 104.
- Setmire, J. [et al.] ; 1993. Detailed study of water quality, bottom sediment, and biota associated with irrigation drainage in the Salton Sea area, California, 1988-90. Prepared in cooperation with the California Regional Water Quality Control Board, Colorado River Basin Region. Sacramento, Calif. : U.S. Geological Survey; 93-4014, 102 p.
- Setmire, J.; Stroud, R. 1990. Reconnaissance investigation of water quality, bottom sediment, and biota associated with irrigation drainage in the Salton Sea area, California, 1986-87. [Sacramento, Calif.] : Dept. of the Interior, U.S. Geological Survey ; 89-4102, 68 p.

- Severson, R.C., Wilson, S.A., and McNeal, J.M., 1987, Analyses of bottom material collected at nine areas in the Western United States for the DOI irrigation drainage task group: U.S. Geological Survey Open-File Report 87-490, 24 p.
- Shacklette, H. and Boerngen, J. 1984. Element concentrations in soils and other surficial materials of the conterminous United States: U.S. Geological Survey Professional Paper 1270, 105 p.
- Stephen, M. 1972. Sedimentary aspects of the New River Delta, Salton Sea, Imperial County, California. Univ. of Southern Calif.
- Stephen, Michael F., and Donn S. Gorsline. 1975. Sedimentary aspects of the New River Delta, Salton Sea, Imperial County, California. Deltas, models for exploration. Broussard, Martha Lou, ed. [Houston]Houston Geological Society. pp. 267-282.
- Stewart, K.; Fey, D.; Hageman, P.; Kennedy, K.; Love, A.; McGregor, R.; Papp, C.; Peacock, T; Sharkey, J.; Vaughn, R.; and Welsch, E. 1992. Department of the Interior, U.S. Geological Survey. Results of chemical analysis for sediments from Department of the Interior National Irrigation Water Quality Program Studies, 1988-1990. Open-File Report 92443. U.S. Geological Survey, Denver Federal Center, Box 25046, MS 973, Denver, CO 80225.
- Thornton, E. and Seyfried, W. Jr. 1975. Sediment-sea-water interaction at 200 and 300' SUP o'C, 500 bars pressure: the role of sediment composition in diagenesis and low-grade metamorphism of marine clay. Geological Society of America Bulletin. pp. 1287-1295.
- Walker, J. 1996. Fresh-Water Inflow and Distribution of Laminated Sediments in the Salton Sea, California. Loma Linda University.
- United States Department of the Interior. 1970. Federal Water Quality Administration. Southwest Regional Office. Salton Sea, California. Water Quality and Ecological Management Considerations. NTIS Accession # PB - 253 691/0/XAB, 54 pp. San Francisco, United States.
- Van de Kamp, P. 1973. Holocene continental sedimentation in the Salton Basin, California: a reconnaissance. Geological Society of America Bulletin. Shell Oil Company Investigation. Vol. 84, pp. 827-48.
- Vonder Haar, S. and Cruz, I. 1979. Fault intersections and hybrid transform faults in the southern Salton Trough geothermal area, Baja California, Mexico. Expanding the geothermal frontier / Geothermal Resources Council Annual Meeting, 24-27 September 1979, Reno, Nevada. pp. 761-764.

Table 1: Concentrations of Inorganic Chemicals in Sediment from the Salton Sea and Surrounding Tributaries Determined to be of Concern

Location	Chemical (concentrations in mg/kg)													
	Antimony	Arsenic	Barium	Cadmium	Chromium	Copper	Molybdenum	Nickel	Selenium	Thallium	Thorium	Uranium	Vanadium	Zinc
Max. Baseline Value mg/kg (a)		22	1,700		200	90	4	66	1.4		20	5.3	270	180
Salton Sea median conc. (mg/kg) (b)		5.6	550		58	28		25	0.7		10.6	4.9	77	78
Whitewater River upstream from HWY 111 (b)		2.4	690	<2	81	34	<2	30	0.1		56	14.6	140	110
Whitewater River at outlet (b)		5	710	<2	210	64	3	170	0.5		18.9	5.5	130	510
Alamo River at international boundary (b)		6.3	510	<2	58	26	<2	26	1.6		12.2	4.8	77	97
Trifolium Drain 1 (b)		5.8	550	<2	53	28	<2	24	1.9		9	4.4	72	78
Ave 64 Evacuation Channel at HWY 195 (b)		4.4	620	<2	75	61	2	2	0.4		21.3	5.1	120	130
New River at midpoint (08/1/86, 08/14/86) (b)		5.4, 11.0	580, 780	<2, <2	63, 73	30, 27	<2, 2	25, 35	0.6, 1.3		10.6, 12.0	6.1, 7.5	77, 96	75, 120
New River at outlet (b)		4.7	720	<2	70	23	<2	22	0.6		19.2	7.7	82	71
East Highline Canal (b)		4.5	690	<2	50	23	<2	22	0.9		12.7	5.9	60	70
Alamo River delta (c)									0.2 - 2.5					
Shoreline Disposal Area (d)		0.9	315		33.9	68.7				0.31			2.6	8.6
Offshore aerobic/anaerobic marine target SSTB (d)	9.9	27.4		1.6			14.5		8.4			14.2	52.5	
Imhoff Tank (d)										0.26				

NOTE:

(a) Shacklette & Boerngen, 1984

(b) Setmire & Stroud, 1990

(c) Selmire [et al.], 1993

(d) Bechtel, 1997 (maximum concentrations reported)

Table 2: Concentrations of Organic Chemicals in Sediment from the Salton Sea and Surrounding Tributaries Determined to be of Concern

Location	Chemical (concentration in µg/kg)															
	Acetone	Carbon disulfide	Chlordane	DBT	DBD	DDE	Dieldrin	Ethylbenzene	gamma-chlordane	Heptachlor	Methoxychlor	PAHs	PCBS	Toluene	Toxaphene	Xylenes
Whitewater River upstream from HWY 111 (b)			<1.0		<0.1	0.6					<0.1		<1		10	
Alamo River outlet (b)			<1.0		20	64					<0.1		<1		<10	
Alamo River at International boundary (b)			<1.0		2.3	18					<0.1		9		<10	
Trifolium Drain 1 (b)			<1.0		3.7	41					<0.1		<1		<10	
Trifolium Drain 1 (e)						110										
Trifolium Drain 4 (b)			<1.0		12	56					<0.1		<1		40	
Vail Drain 4 (b)			<1.0		7.8	57					45		<1		<10	
Ave 64 Evacuation Channel at HWY 195 (b)			1		5.8	56					<0.1		<1		<10	
Ave 64 Evacuation Channel at HWY 195 (e)						67										
New River at midpoint (08/14/86) (b)			5		3.5	7.4					<0.1		4		<10	
New River at international boundary (b)			20		24	7.6					<0.1		24		<10	
East Highline Canal (b)			<1.0		2.3	18					<0.1		9		<10	

Table 2: Concentrations of Organic Chemicals in Sediment from the Salton Sea and Surrounding Tributaries Determined to be of Concern


Location	Chemical (concentration in ug/kg)															
	Acetone	Carbon disulfide	Chlordane	DDT	DDD	DDF	Dieldrin	Ethylbenzene	gamma-Chlordane	Heptachlor	Methoxychlor	PAHs*	PCBs	Toluene	toxaphene	Xylenes
Shoreline Disposal Area (d)	23	2		3.1	4.9	6.6	3	2	3.4	3.5	14	85		15		11
Imhoff Tank (d)						3.2	0.6		190	290						
1 mile from Whitewater River outlet (f)	0-11.5 cm			<25	5	5	<5									
	11.5-23 cm			<25	<5	<5	<5									
2.5 miles from Whitewater River outlet (f)	0-11.5 cm			<25	5	5	<5									
	11.5-23 cm			25	20	23	<5									
5 miles from Whitewater River (f)	0-11.5 cm			<25	12	14	<5									
	11.5-23 cm			25	5	5	5									
1 mile from Alamo River outlet (f)	0-11.5 cm			25	5	5	92									
	11.5-23 cm			25	5	5	100									
2.5 miles from Alamo River outlet (f)	0-11.5 cm			25	5	16	49									
	11.5-23 cm			82	5	18	880									
5 miles from Alamo River outlet (f)	0-11.5 cm			25	5	5	60									
	11.5-23 cm			25	5	5	43									

NOTE:

- (a) Shacklette & Boerngen, 1984
- (b) Setmire & Stroud, 1990
- (c) Setmire [et al.], 1993

- (d) Bechtel, 1997 (maximum data reported)
- (e) Eccles, 1979
- (f) Hogg, 1973

* Polycyclic Aromatic Hydrocarbon (PAHs) values are for Benzo(a)anthracene and Chrysene



Avenue of Erosion Channel at Highway 195

EXPLANATION

- Salton Assessment Area
Setmire (et al), 1993.
- Sediment Investigation
Setmire and Stroud, 1990.
- Sediment Grain Size
 - Sand 250-1,000 µm
 - Fine Sand 64-250 µm
 - Fine Silt 16-64 µm
 - Silt 4-16 µm
 - Clay < 4 µm

NOTE: Grain Size of lake bottom sediments (after Arnel, 1961)



0 6 Miles



64-24-11 1200000000

Salton Sea
Salton Sea
Bottom Sediment Sampling Sites
Figure 1
Project No. 6624

MAP SOURCES:

- State of California - Division of Mines and Geology, Geologic Map of California (Jenkins), Salton Sea sheet, 1977.
- U.S.G.S Topographic Map, 1:250,000, Santa Ana, California, 1961.